arXiv:astro-ph/0405125v1 6 May 2004

Extragalactic Ultracompact HII Regions: Probing the Birth Environments of Super Star Clusters

Kelsey E. Johnson

Astronomy Dept., University of Wisconsin, 475 N. Charter St., Madison, WI, U.S.A.

Abstract. In recent years, a number of extragalactic massive star clusters that are still deeply embedded in their birth material have been discovered. These objects represent the youngest stage of massive star cluster evolution yet observed, and the most massive and dense of these may be proto globular clusters. Their properties appear to be similar to those of ultracompact HII regions in the Galaxy, but scaled up in total mass and luminosity. In many cases, these clusters are only visible at mid-IR to radio wavelengths, and they have typically been detected as "inverted" spectrum radio sources. However, the set of existing observations is anemic, and our current physical model for these natal clusters in simplistic. This article will overview what we think we know about these objects based on existing observations and outline some of the most significant gaps in our current understanding.

1. Introduction

There is no question that massive star clusters, and super star clusters in particular, are among the most extreme star formation environments in the local universe. Moreover, these clusters can have a dramatic affect on their surrounding interstellar medium, and in some cases the intergalactic medium. Nevertheless, the conditions required for their formation remain far from understood. As just one example, there is still significant debate over whether super star clusters (SSCs) are merely the result of populating a cluster initial mass function statistically, or whether they truly require a separate mode of star formation. While a multitude of adolescent massive star clusters have been observed at optical wavelengths, these observations cannot penetrate the natal material surrounding the youngest clusters in order to reveal the properties of their birth environments (Figure 1). Nevertheless, if we want to understand the formation of massive star clusters, it is not a terribly bad idea to observe them while they are forming.

A number of investigations at infrared (IR) and radio wavelengths have now revealed a population of extremely young massive star clusters in a growing sample of galaxies (e.g. Turner, Ho, & Beck 1998; Kobulnicky & Johnson 1999; Tarchi et al. 2000; Beck, Turner, & Gorjian 2001; Gorjian, Turner, & Beck 2001; Vacca, Johnson, & Conti 2002; Johnson et al. 2001; Neff & Ulvestad 2000; Plante & Sauvage 2002; Beck et al. 2002; Johnson, Indebetouw, & Pisano 2003; Johnson & Kobulnicky 2003). These observations suggest that the early stages of massive star cluster evolution may parallel those of individual massive stars observed in the Milky Way. Therefore, our knowledge about individual ultracompact HII regions (UCHIIs) is critical to understanding and interpreting observations of embedded


Evolutionary State
Observable Regimes
Deeply Embedded Ultra-young Cluster
Mid-IR - Radio (cm)
Highly Extincted Young Cluster
Near-IR - Radio (cm)
Fully Emerged Adolescent Cluster
Optical - Near-IR

Figure 1. A cartoon illustrating stages in the early evolution of massive star clusters. Observations in at thermal infrared to radio wavelengths are required to probe their birth environments.

massive star clusters. Of course, individual massive star formation is also poorly understood, and we would be remiss to blindly extrapolate the inferred properties of individual massive proto-stars to the formation of SSCs. Perhaps the most obvious way in which this extrapolation could lead us astray is by neglecting the effects of the density of proto-stars on star formation. In fact, it seems likely that a tremendous amount of physical insight will come from trying to understand how the properties of star formation scale between the two extremes of *individual* massive star formation and *SSC* formation.

Despite the pioneering observations done by a number of investigators, our current knowledge about embedded massive star clusters is anemic, and we have only begun to investigate their properties, environments, and evolution. In the pages that follow, I will discuss the current observational limitations (§2.), give a brief tour of the sample of extragalactic embedded clusters (§3.), overview what we think we know about these objects (as well as what we do not know, §4.), and outline the impact that the next generation of powerful observatories at infrared to radio wavelengths will have on this area of research (§5.).

2. Observational Limits

Because star formation takes place within dense and dusty cocoons, we can only probe these regions with observations at infrared or longer wavelengths. In fact, some natal clusters (presumably the youngest) are even invisible at near-IR wavelengths as long as the K-band ($\sim 2.2\mu$m, Vacca, Johnson, & Conti 2002) or even the L-band ($\sim 3.8\mu$m, M. Sauvage, private communication). On the other hand, radio emission is completely unfettered by dust, but the observations need to be sensitive at high frequencies (> 5 GHz) in order to avoid self-absorption by



Figure 2. The 5σ sensitivities for 8 hr integrations at 6 cm for various radio interferometers (both current and planned) are indicated by the horizontal grey shaded areas. The flux densities of "typical" UCHIIs and natal super star clusters as a function of distance are shown with the diagonal lines. The distances to a handful of reference galaxies are indicated.

the dense gas and disentangle non-thermal radio emission. Of course, obtaining the necessary sensitivity is only half of the battle; achieving adequate spatial resolution is also a limiting constraint with current facilities. Roughly speaking, a linear resolution of at least ~ 0.5 pc is necessary to observe individual UCHIIs, and embedded SSCs require a linear resolution of ~ 10 pc or better.

In principle, mid-IR to sub-millimeter wavelengths are ideal for observing extragalactic UCHIIs and natal SSCs because these objects are extremely luminous in this regime due to thermal dust emission (Figure 5). As a demonstration of their potency at thermal wavelengths, UCHIIs are among the most luminous Galactic objects in the *IRAS* catalog ($>60\%$ of the *IRAS* sources with flux densities $> 10^4$ Jy at 100μm are UCHIIs, Wood & Churchwell 1989), and the embedded SSCs in the starburst galaxies He 2-10 and NGC 5253 are responsible for nearly all of the mid-IR emission from these galaxies (Beck, Turner, & Gorjian 2001; Gorjian, Turner, & Beck 2001; Vacca, Johnson, & Conti 2002). However, we are severely limited by spatial resolution with the facilities currently available at infrared to sub-millimeter wavelengths. This situation will improve to some degree with the next generation of observatories that are beginning to come online; however, even the *Spitzer, Herschel,* and *SOFIA* telescopes will not obtain spatial resolutions of better than $\sim 1''$ at their shortest wavelengths.

The sensitivity and spatial resolution of existing radio interferometers is somewhat better matched to observing extragalactic UCHIIs and natal SSCs. Figure 2 illustrates the sensitivities of various radio interferometers. The current facilities have the sensitivity to detect individual UCHIIs nearly as far away as the starburst galaxy NGC 253, or embedded SSCs nearly out to ultralu-



Figure 3. Examples of the most nearby and most distant embedded clusters yet studied. **(left)** The HII region N81 in the SMC; an *HST* Hα image is shown in gray-scale, and *ATCA* 3 cm radio emission is shown in contour (Indebetouw, Johnson, & Conti 2004). **(right)** The starburst galaxy SBS0335-052; an *HST* I-band image is shown in gray-scale, and *VLA* 1.3 cm radio emission is shown in contour (Johnson & Plante in prep.)

minous infrared galaxy Arp 220. Perhaps the most obvious places to look for extragalactic UCHIIs are the Magellanic Clouds, however the best spatial resolution available in the southern hemisphere from the *Australia Telescope Compact Array* at this distance is ∼ 1 pc, which limits our ability to asses whether HII regions are actually "ultra" compact. Another obvious place to search of UCHIIs is in M33, the sister galaxy to the Milky Way. However, a linear resolution of better than $\sim 0.1''$ is required in order to observe UCHIIs at this distance (840 kpc), which will be difficult (but not impossible) to achieve with the *Very Large Array* (VLA). The constraints are somewhat more relaxed for observing embedded SSCs, which are both significantly larger and brighter than individual UCHIIs. With the *VLA*, it is possible to resolve embedded SSCs in galaxies out to distances of ∼ 10 Mpc. Beyond this distance, both crowding and contamination by non-thermal and optically-thin thermal radio emission becomes an issue, and one has to use caution when interpreting the data.

3. Sample of Extragalactic Natal Clusters

A number of embedded massive star clusters have been identified in a variety of galaxies. The objects that have been most well-studied to date are those that were identified in the incipient radio observations of the starburst galaxies NGC 5253 (∼ 4 Mpc) and He 2-10 (∼ 9 Mpc) (Turner, Ho, & Beck 1998; Kobulnicky & Johnson 1999). Subsequent work has revealed a number of embedded star forming regions in a range of galactic environments. In the most nearby examples, Indebetouw, Johnson, & Conti (2004) have identified a number of candidate UCHIIs and UCHII complexes in the Magellanic Clouds (here the word "candidate" is used because the spatial resolution is not adequate to determine whether the regions are "ultra" compact). In this sample, the radio observations indicate stellar ionizing fluxes and morphologies consistent with sources ranging from single UCHIIs to relatively modest UCHII associations (Figure 3a).

Compact radio-detected clusters have also been identified in galaxies such as NGC 6822 at ~ 0.5 Mpc, IC 4662 at ~ 2 Mpc, IIZw40 at ~ 11 Mpc, Haro 3 at ~ 13 Mpc, NGC 5398 at ~ 17 Mpc, NGC 2146 at ~ 15 Mpc, the Antennae at ~ 21 Mpc, and SBS 0335-052 at ~ 53 Mpc (Neff & Ulvestad 2000; Tarchi et al. 2000; Beck et al. 2002; Johnson, Indebetouw, & Pisano 2003; Johnson et al. 2004, Johnson & Plante in prep., Seth & Johnson in prep.) [1]. The star forming regions in this sample have properties ranging from those of natal OB-associations to incredibly massive SSCs. For example, the main radio source in SBS 0335-052 has an ionizing flux equivalent to $\sim 12,000$ O7-type stars (Figure 3b; note that the linear resolution of these observations is only ~ 75 pc, and this source is likely to be composed of several individual clusters).

4. What We Think We Know & What We Know We Don't Know

Given the limited observations available at the present time, attempts to constrain the physical properties of embedded clusters have been fairly crude. As a result, the current physical model we have for these objects is simplistic: When an SSC is born, the massive stars ionize the surrounding natal material, creating an extremely dense HII region. The dense HII region is in turn surrounded by a dust cocoon that is presumably quite warm on the inner boundary, with the temperature decreasing at distances farther from the ionizing stars. The constituent ionizing stars may be surrounded by individual HII regions and dust cocoons in the earliest stages, but this is currently unknown. Clusters that are more sparse are likely to resemble ultracompact HII region (UCHII) complexes in the Milky Way (such as W49A), and contain spatially discrete HII regions and cocoons, but the existence of individual cocoons becomes less tenable in the extreme stellar densities of the cores of massive SSCs.

The dense HII regions are characterized by an "inverted" radio spectral energy distributions due to optically thick free-free emission ($\alpha > 0$, where $S_\nu \propto \nu^\alpha$). The frequency at which the radio emission becomes optically thick is largely dependent on the density of the HII region. In the earliest stages of cluster evolution, all of the stellar luminosity is reprocessed by the surrounding dust cocoon, and this cocoon is observable at thermal infrared to sub-millimeter wavelengths. To some extent, the nature of the resulting spectral energy distribution depends on whether the dust in the cocoon is smooth or clumpy, but this is not currently constrained by observations.

Existing observations have provided *estimates* for some of the physical properties of natal clusters, including size, electron density, pressure, stellar mass, HII mass, dust mass, and age. For the most nearby clusters, their size can be directly measured using high spatial resolution observations. For more distant clusters, model HII regions can be used to fit the radio spectral energy distribution and infer both the radius and electron density (Figure 4b). The

---

[1] This list in not intended to be exclusive or exhaustive.



Figure 4. **(left)** Natal SSCs in the starburst galaxy He 2-10. An *HST* I-band image is shown in gray-scale, and *VLA* 2cm emission is shown in contour. The four radio sources without optical counterparts are natal SSCs. The arrow indicates the source for which the spectral energy distribution is shown. **(right)** The radio fluxes for the most luminous natal SSC in He 2-10 along with the best fit spectral energy distribution for a model HII region.

embedded SSCs that have been analyzed in this way (e.g. those in He 2-10, Johnson & Kobulnicky 2003) have HII region radii of only a few parsecs, and the electron densities have global values of $\sim 10^3$ cm^{-3}, with peak values in excess of $\sim 10^6$ cm^{-3}. Assuming temperatures of $\sim 10^4$ K, these densities imply global pressures of $P/k_B > 10^7$ K cm^{-3} and peak pressures reaching values of $P/k_B \sim 10^{10}$ K cm^{-3}.

The masses associated with the stellar clusters, HII regions, and dust cocoons can be bootstrapped from radio and infrared observations. The optically-thin thermal flux (typically measured with the highest available radio frequency) can be used to infer a total ionizing luminosity, and the ionizing luminosity can be translated into an embedded stellar mass by assuming a stellar initial mass function. Using this technique, clusters with a range of masses have been detected (spanning modest OB-associations to massive SSCs). The HII mass can be estimated given the radii and densities for the ionized regions. In at least some cases, the HII mass has anomalously low values ($< 5\%$ of the stellar mass, Johnson & Kobulnicky 2003), which we tentatively interpret as a sign of the youth of the HII regions. The mass of the dust cocoons around embedded SSCs has also been estimated in a few cases using infrared observations; these estimates have suggested dust masses of $\sim 10^5 - 10^6 M_\odot$ for massive SSCs in SBS0335-052 and He 2-10 (Plante & Sauvage 2002; Vacca, Johnson, & Conti 2002).

The ages of the natal clusters are exceptionally tricky to ascertain, as there are no available spectral diagnostics to determine age at wavelengths longer than the near-IR (and for shorter wavelengths the diagnostics are unreliable for ages less than ~ 1 Myr in any case). The best we can do is offer circumstantial evidence and look at the back of a proverbial envelope. There are two independent "back-of-the-envelope" calculations that can be used to constrain the lifetimes of the embedded phase of SSC evolution: (1) The pressures inferred for these regions are several orders of magnitude higher than those typical in the

ISM. As a result, the HII regions ought to expand and dissipate on very short time-scales ($\tau < 10^6$ yrs). However, there are caveats to this argument: First, we do not know the pressure of the surrounding ISM, and it could be a good deal higher than normal since it is a starburst region. Second, if the clusters are massive enough, the gravity could be large enough to help balance pressure and confine the HII region (e.g. Turner et al. 2003). (2) If star formation has been relatively continuous over ~ 10 Myr, then a comparison between the total number of clusters with ages < 10 Myr and the number embedded clusters ought to reflect the their relative lifetimes. When this has been applied to galaxies with good enough statistics (e.g. He 2-10, Kobulnicky & Johnson 1999), the clusters appear to spend $\sim 10-20\%$ of the lifetimes of the constituent massive stars in the embedded phase ($\tau \sim 0.5-1$ Myr). There are caveats to this argument as well. First, it relies on star formation being continuous over the lifetimes of massive stars. Second, we have to make sure that both the optical and radio observations are sensitive to roughly the same cluster masses. In any case, it is comforting that both calculations suggest the same lifetimes for the embedded phase of cluster evolution, so our estimates are probably not too far off.

While we have estimates for a few of the quantities associated with the birth of massive star clusters, there are a potpourri of properties that it would be nice to know more about. For example, it would be instructive to *directly* measure densities, pressures, and temperatures using various line diagnostics at IR to radio wavelengths. Another quantity that might be particularly useful for theorists (that is almost completely unconstrained) is the star formation efficiency in these objects. We also have no idea how much ionizing radiation could be escaping from the natal cocoons (and it could be quite a lot if the dust is clumpy). In fact, we know almost nothing about the natal dust cocoons (including the temperatures, grain compositions, and geometry). This list is certainly not exhaustive. An understanding of these quantities, and how they relate to the properties of the resulting cluster, will eventually provide a great deal of insight into the requirements for massive star cluster formation.

5. Future Prospects

While there are no lack of questions remaining about the birth of massive star clusters, it is an exceptional good time to be asking these questions. Over the next decade (or so), a range of powerful facilities are going to become available at infrared to radio wavelengths. Figure 5 illustrates the spatial resolution and sensitivity of the IR to millimeter facilities that will soon become available. In the thermal infrared, *Spitzer, Herschel*, and *SOFIA* will provide a vast improvement over previous facilities, and embedded star forming regions are likely to be a dominate source of flux in the thermal spectral energy distribution of star forming galaxies. However, we will have to be cautious in our interpretation of data from these observatories because they will not have the spatial resolution necessary to isolate individual natal super star clusters. At millimeter wavelengths, the *Atacama Large Millimeter Array* will open up an entirely new window for observing extragalactic star formation with resolution and sensitivity incredibly well-matched to observing embedded clusters. In the radio regime (see Figure 2), the *Expanded Very Large Array* and *Square Kilometer Array* are



Figure 5. **(left)** The anticipated spatial resolution and wavelength coverage for upcoming infrared and millimeter observatories. **(right)** The sensitivities of these observatories normalized to 500s integration time. A model spectral energy distribution (Whitney et al. in prep) for a $10^6 M_\odot$ natal SSC at a distance of 10 Mpc is over-plotted with a dashed line for reference.

going to achieve sensitivities and resolutions that will allow us to study nearby natal clusters in fantastic detail and enable us to discover new natal clusters out to distances of 100 Mpc and beyond.

Acknowledgments. This work has benefited from many fruitful discussions with my collaborators, including P.Conti, M.Goss, R.Indebetouw, H.Kobulnicky, and W.Vacca. I also gratefully acknowledge support for this work provided by the NSF through an Astronomy and Astrophysics Postdoctoral Fellowship. Support for proposal #09934 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

References

Beck, S.C., Turner, J.L., Langland-Shula, L.E., Meier, D.S., Crosthwaite, L.P., Gorjian, V. 2002, AJ 124, 2516
Beck, S.C., Turner, J.L., & Gorjian, V. 2001, AJ 122, 1365
Gorjian, V., Turner, J.L., & Beck, S.C. 2001, ApJ 554
Indebetouw, R., Johnson, K.E., & Conti, P.S. 2004, AJ submitted
Johnson, K.E., Indebetouw, R., Watson, C., & Kobulnicky, H.A. 2004, AJ, submitted
Johnson, K.E. & Kobulnicky, H.A. 2003, ApJ 597, 923
Johnson, K.E., Indebetouw, R., & Pisano, D.J. 2003, AJ 126, 101
Johnson, K.E., Kobulnicky, H.A., Massey, P., & Conti, P.S. 2001, ApJ 559, 864
Kobulnicky, H.A. & Johnson, K.E. 1999, ApJ 527, 154
Plante, S. & Sauvage, M. 2002, AJ 124, 1995
Neff, S.G. & Ulvestad, J.S. 2000, AJ 120, 670
Tarchi, A., Neininger, N., Greve, A., Klein, U., Garrington, S.T., Muxlow, T.W.B., Pedlar, A., Glendenning, B.E. 2000, A&A 358, 95
Turner, J.L., Beck, S.C., Crosthwaite, L.P., Larkin, J.E., McLean, I.S., Meier, D.S. 2003, Nature, 423, 621
Turner, J.L., Ho, P.T.P., & Beck, S.C. 1998, AJ 116, 1212
Vacca, W.D., Johnson, K.E., & Conti, P.S. 2002, AJ 123, 772
Wood, D.O. & Churchwell, E. 1989, ApJS 69, 831